

PULSin

in vitro Protein, Antibody and Peptide Delivery reagent

PROTOCOL

DESCRIPTION

PULSin is a powerful reagent dedicated to the delivery of peptides, antibodies and proteins into cells. It contains a cationic amphiphile molecule whose formulation is proprietary. PULSin delivers anionic proteins and antibodies to a large variety of eukaryotic cell lines including primary cells. PULSin is most efficient when able to interact with the protein by electrostatic and/or lipophilic interactions. Thus, anionic proteins (i.e. proteins with an isoelectric point < 7) and antibodies are particularly well-suited for delivery with PULSin. Yet delivery is not restricted to anions, as most proteins have a lipophilic core.

1	Delivery protocol for adherent cells	2
1.1	Cell seeding	2
1.2	Protein Delivery Protocol.....	2
1.3	Protein delivery protocol for sensitive cells	3
1.4	Optimization guidelines.....	4
2	Delivery protocol for suspension cells	6
2.1	Cell seeding	6
2.2	Protein delivery protocol	6
3	Troubleshooting	7
4	Product Information.....	8

1 DELIVERY PROTOCOL FOR ADHERENT CELLS

1.1 CELL SEEDING

For optimal protein delivery with PULSin, the cells should be 70-80% confluent on the day of the experiment. Typically, for protein delivery in 24-well plates, 70 000 to 100 000 cells are seeded per well and incubated for 16 to 24 h. Refer to Table 1 for protein delivery in other culture formats. Since the efficiency of protein delivery partly depends on cell confluency, the culture density should be optimized for each cell type.

Table 1. Recommended number of cells to seed the day before protein delivery with PULSin

Culture vessel	Number of adherent cells to seed	Surface area per well (cm ²)	Volume of medium per well to seed the cells (ml)
96-well	10 000 - 15 000	0.3	0.2
24-well	70 000 - 100 000	1.9	1
12-well	100 000 - 180 000	3.8	2
6-well / 35 mm	200 000 - 300 000	9.4	3
60 mm / flask 25 cm ²	300 000 - 800 000	25 - 28	5
100 mm / flask 75 cm ²	1.10 ⁶ - 2.10 ⁶	75 - 78.5	10

1.2 PROTEIN DELIVERY PROTOCOL

The following protocol is given per well of a 24-well plate, for the delivery of 1 µg of protein, antibody or peptide.

- For R-PE, use 4 µl of PULSin per µg of protein.
- For antibodies, start with 2.5 µl of PULSin per µg of antibody.

For other culture format, please refer to Table 2. Guidelines for optimization are given in section 1.3.

1. Dilute 1 µg of protein in 100 µl of 20 mM HEPES in a microcentrifuge tube. Vortex gently and spin down briefly.
2. Vortex PULSin reagent for 5 sec and spin down before use.
3. Add 4 µl of PULSin. Vortex immediately and spin down briefly.
4. Incubate for 15 minutes at room temperature.
5. Wash cells once with 1X PBS or culture medium without serum. The washing step is critical to remove all traces of serum.
6. Add 900 µl of culture medium without serum per well.

7. Add 100 μ l of protein/PULSin mix per well and homogenize by gently swirling the plate.
8. After 4 hours of incubation at 37°C, remove the medium containing the protein/PULSin complexes and replace with complete growth medium. Return the plate to the incubator.
9. Analyze protein activity or visualize intracellular fluorescence immediately or after an incubation period.

N.B.: For R-PE, the excitation by a laser at 488 nm induces a maximal light emission at 575 nm, and the optimal time of visualization is 16 hours after protein delivery.

Table 2. Protein delivery guidelines according to the cell culture vessel

Culture Vessel	Amount of protein (μ g)	Volume of 20 mM Hepes Buffer (μ l)	Volume of PULSin (μ l)
96-well	0.3	20	1.2
24-well	1	100	4
12-well	2	150	8
6-well / 35 mm	4	200	16
60 mm / flask 25 cm ²	7	400	28
100 mm / flask 75 cm ²	10	1000	40

1.3 PROTEIN DELIVERY PROTOCOL FOR SENSITIVE CELLS

Sensitive cells may not withstand the absence of serum for 4 hours, thus we suggest an alternative protocol below. Proceed with Steps 1 to 6 as in 1.2, then add a short centrifugation step as indicated below.

1. Gently centrifuge the cells for 5 min at 190 g (if the cells can withstand it) directly after adding the protein/PULSin complexes onto the cells.
2. Incubate for 30 min at 37°C. Remove the medium containing the protein/PULSin complexes and replace with complete growth medium. Return the plate to the incubator.

Analyze protein activity or visualize intracellular fluorescence immediately or after an incubation period.

1.4 OPTIMIZATION GUIDELINES

Optimization is highly recommended to obtain as high protein delivery efficiency as possible with PULSin. In addition, variations may be observed from one cell line to another, even with the same protein. Thus we recommend testing a range from 0.5 μg to 4 μg of protein, antibody or peptide and 1 μl to 4 μl of PULSin per well of 24-well plate (cf. Table 3).

Table 3. Optimization guidelines for PULSin-mediated protein delivery (per well of 24-well plate)

Amount of protein (μg)	0.5			2			4		
Volume of PULSin (μl)	1	2	4	1	2	4	1	2	4


The amounts to be tested are the same for proteins, antibodies or peptides.

Browse our cell transfection database to find the optimized conditions according to your cell line.

<http://www.polyplus-transfection.com/resources/cell-transfection-database/>



Protein delivery protocol for a well of 24-well plates



2 DELIVERY PROTOCOL FOR SUSPENSION CELLS

2.1 CELL SEEDING

For protein delivery in suspension cells (Jurkat, THP-1, K-562, etc.), cells are counted and collected by centrifugation (190g for 5 minutes) on the day of the experiment. Per well of a 24-well plate, seed 5×10^5 cells in 1 ml of Opti-MEM[®] (without serum but containing glutamine). The following protocol is given as a starting point, for optimization please refer to section 1.3.

2.2 PROTEIN DELIVERY PROTOCOL

100 μ l of protein/PULSin complexes are required per well of 24-well plate and are prepared as follows:

1. Dilute 1 - 2 μ g of protein into 100 μ l of 20 mM Hepes. Vortex gently and spin down briefly.
2. Vortex PULSin reagent for 5 sec and spin down before use.
3. Add 2 - 4 μ l of PULSin into each tube. Vortex immediately and spin down briefly.
4. Incubate for 15 minutes at room temperature and add the protein/PULSin mix into the 1 ml of cells at the density of 5×10^5 cells/ml.
5. After 4 hours of incubation at 37°C, centrifuge the cells 5 min at 190g and resuspend them in 1 ml complete growth medium. Return the plate to the incubator.
6. Analyze protein activity or visualize intracellular fluorescence immediately or after an incubation period

3 TROUBLESHOOTING

Observations	Actions
Low protein delivery efficiency	Ensure that adherent cells are 70-80% confluent on the day of the experiment.
	Optimize the amount of protein delivered (0.5 to 2 µg per well of 24-well plate).
	Include an additional washing step with PBS to ensure that all traces of serum have been removed.
	Use protein as pure as possible.
	Optimize the protein/PULSin ratio from 1:1 to 1:4.
	Perform a positive control delivery experiment with the positive control protein R-phycoerythrin (included in each PULSin kit), using 1 µg per well of 24-well plate on your cells.
	Ensure that the complexes are prepared in HEPES buffer and added to the cells in serum-free medium.
Presence of aggregates	Ensure that cells are more confluent on the day of delivery, ideally 80%.
Cellular toxicity	Reduce the amount of protein used in the assay.
	Check protein concentration and ensure that the protein /PULSin ratio is lower than 1:4.
	Reduce the incubation time of the protein/PULSin complexes with the cells from 4 h to 2 h. For very sensitive cells, include a centrifugation step (5 min at 190g right after adding the protein/PULSin complexes to the cells) and incubate only 30 min as described in section 1.4.

4 PRODUCT INFORMATION

4.1 ORDERING INFORMATION

Ref #	PULSin Reagent	Number of delivery experiments
501-01	0.1 ml	6 delivery experiments in 6-well plate
501-04	0.4 ml	25 delivery experiments in 6-well plate

4.2 CONTENT

- 0.4 ml of PULSin reagent is sufficient to perform ca. 100 experiments in 24-well plates or ca. 25 experiments in 6-well plates.
- R-phycoerythrin (R-PE) (20 µg) to be used as a positive control at 0.1 µg/µl. The excitation of R-PE by 488 nm laser light induces a maximal light emission at 575 nm.
- HEPES Buffer (20 mM), 20 ml (ref # 501-01 and 501-04) for protein dilution.

4.3 REAGENT USE AND LIMITATIONS

For research use only. Not for use in humans.

4.4 QUALITY CONTROL

Every batch of PULSin is tested by delivering R-phycoerythrin into HeLa cells.

4.5 FORMULATION AND STORAGE

PULSin is provided as an aqueous solution in sterile and apyrogenic water. PULSin, R-phycoerythrin and HEPES buffer are shipped at 4°C, should be stored at 4°C upon arrival, and as guaranteed by the Certificate of Analysis, will be stable for at least one year when stored appropriately.

Polyplus-transfection® has been awarded ISO 9001 Quality Management System Certification since 2002, which ensures that the company has established reliable and effective processes for manufacturing, quality control, distribution and customer support.

4.6 TRADEMARKS

Polyplus-transfection is a registered trademark of Polyplus-transfection.

How to cite us: "PULSin (Polyplus-transfection S.A, Illkirch, France)"

4.7 TECHNICAL ASSISTANCE AND SCIENTIFIC ADVICE

Contact the friendly Polyplus technical support *via*:

- The Polyplus website: www.polyplus-transfection.com
- Email: support@polyplus-transfection.com
- PHONE: + 33 (0)3 90 40 61 87